arXiv:1304.3925v1 [quant-ph] 14 Apr 2013

Long-range entanglement is necessary for a topological storage of quantum information

Isaac H. Kim[1]

[1]*Institute of Quantum Information, California Institute of Technology, Pasadena CA 91125, USA*

(Dated: April 16, 2013)

We ask whether the ground state entanglement of a quantum many-body system imposes a fundamental constraint on its information storage capacity. We show that the amount of long-range entanglement gives a rigorous upper bound to this problem, even without invoking the properties of the parent Hamiltonian. In particular, we show that $\log N \leq 2\gamma$ for a large class of topologically ordered systems on a torus, where N is the number of topologically protected states and γ is the topological entanglement entropy. We also obtain a necessary condition for a quantum error correcting code to have a nonvanishing rate: its average entanglement entropy over the subsystems smaller than the code distance must satisfy a strict volume law.

Entanglement entropy is a canonical measure for quantifying entanglement in a bipartite pure state.[1] There has been a recent surge of interest in studying entanglement entropy in a ground state of a quantum many-body system.[2] One of the motivations behind these studies is that the entanglement entropy is a useful probe for detecting the phase of the quantum many-body system. For example, entanglement entropy in one dimensional critical systems follows a universal logarithmic scaling law, and its prefactor is related to the conformal charge of the theory.[3] In two spatial dimensions, the quantum dimension of the topological quantum field theory describing the low-energy physics can be inferred from a constant subcorrection term of the entanglement entropy.[4–6]

Another important motivation comes from the numerical simulation of quantum many-body systems. Classes of variational ansatz such as the matrix product states(MPS),[7, 8] projected entangled pair states(PEPS),[9] and the multi-scale entanglement renormalization ansatz(MERA)[10] have certain entropy scaling laws. Since these variational states reproduce the entanglement scaling of gapped/critical systems, they are suitable for efficiently simulating the ground state properties of the quantum many-body systems. In particular, a rigorous argument can be made for one-dimensional gapped systems, where an explicit scaling relation between the entanglement entropy and the MPS bond dimension is known.[11–13]

Recently another possibility has been explored by several authors.[14–16] Their approaches differ from the conventional ones in several aspects. For one thing, they explicitly use the entanglement entropy over multiple subsystems simultaneously. Moreover, the main objective of these works is not necessarily focused on obtaining an order parameter of the phase. Rather, they are interested in identifying a hidden structure of the quantum many-body phase that protects their universal properties.

This hidden structure by in large can be attributed to the structure of states that are approximately conditionally independent. A tripartite state ρ_{ABC} is conditionally independent if its conditional mutual information $I(A:C|B) = S(AB) + S(BC) - S(B) - S(ABC)$ is equal to 0, where $S(A) = -\mathrm{Tr}(\rho_A \log \rho_A)$ is the entanglement entropy of the subsystem A. Similarly, a tripartite state is approximately conditionally independent if its conditional mutual information has a nonzero value that is close to 0. At least for gapped quantum many-body systems, there is a good reason to expect such subsystems to appear quite naturally.[14] The power of these approaches lies on the fact that their argument only relies on the generic properties of the ground state alone. Moreover, the structure that arises from the vanishing conditional mutual information is manifestly nonlocal: it gives a nontrivial constraint between the density matrices over large regions.[17–19]

In this Letter, we explore this structure further in the context of finding a fundamental limit on the information storage capacity of a quantum many-body system. There has been a number of results in recent years, where information storage tradeoff bounds for *local quantum codes* have been obtained.[20–24] A local quantum code refers to a set of degenerate ground states stabilized by a sum of geometrically local commuting projectors. Important examples include quantum double and Levin-Wen model.[25, 26] Bravyi *et al.* showed that the following bound holds for such systems

$$kd^{2/(D-1)} = O(n), \tag{1}$$

where k is the number of qubits, d is the code distance, D is the spatial dimension of the lattice, and n is the number of particles.[21]

Local quantum codes cover a rich array of systems. Indeed, it has been conjectured that Levin-Wen model can describe any non-chiral topologically ordered systems in two spatial dimensions.[26] However, none of these models give rise to the chiral gapless boundary excitations. Hence, the tradeoff bound cannot be applied to physical systems exhibiting the integer quantum Hall effect or the fractional quantum Hall effect.[27, 28] Furthermore, there are models that are topologically ordered yet do not necessarily have a local commuting parent hamiltonian. Examples include certain quantum dimer models,[29, 30] parent hamiltonian of PEPS,[31, 32] and Kitaev's honeycomb model.[33]

To deal with such systems, we approach this problem by focusing only on the properties of the state. More precisely, we obtain an upper bound on the number of topologically protected qubits in terms of a linear combination of entanglement entropies over some local regions of a *single wavefunction.* Entanglement entropies are later related to the code parameters. At first, the preceding claim might seem oxymoron to some of the readers. How can one bound a number of degenerate ground states without knowing the parent hamiltonian? Further, how can one even define an entanglement entropy of a subsystem if there are multiple states? Answers to both of these questions lie on an important difference between a classical error correcting code and a quantum error correcting code.

The key property of the quantum error correcting code that sets apart from its classical counterpart is the local indistinguishability property. While a more general version of this definition exists,[34–36] it will be convenient to use the concept of topological quantum order(TQO)[37] for our purpose. Following Ref.37, we say that a set of states $\{|\psi_i\rangle\}_{i=1,\cdots,N}$ satisfies a TQO condition with (r,ϵ)-error if

$$\begin{aligned} |\langle\psi_i|\,\phi\,|\psi_j\rangle| &\leq \|\phi\|\epsilon, \quad i\neq j \\ |\langle\psi_i|\,\phi\,|\psi_i\rangle - \langle\psi_j|\,\phi\,|\psi_j\rangle| &\leq \epsilon\|\phi\|, \end{aligned} \tag{2}$$

holds for any ϕ that is restricted to a ball of radius r, where $\|\cdots\|$ is the operator norm. If the approximation parameters r and ϵ are obvious from the context, we shall simply say that the states are locally indistinguishable. An important consequence of Eq.2 is that the reduced density matrices of the locally indistinguishable states are close to each other in a trace distance. Therefore, one can unambiguously define the entanglement entropy of the aforementioned set of states up to a small error, so long as the subsystem can be contained in a ball of radius r. We shall call such subsystems to be *local.*[59]

Roughly speaking, each of the qubits in the quantum error correcting code is entangled to each other so that the information can be stored nonlocally. As the number of encoded qubits increase, more entanglement is necessary to distribute the information sufficiently nonlocally. Hence, there has to be an inherent limit on the number of protected qubits if the entanglement across different subsystems are bounded.

In this Letter, we shall set this intuition on a rigorous ground. Our approach is purely information-theoretic: we only use the strong subadditivity of entropy as our main technical tool.[38] However, we believe it would be more instructive to discuss some of the related results to understand the motivation behind our construction. Strong subadditivity of entropy asserts that the conditional mutual information $I(A:C|B)$ of a tripartite quantum state ρ_{ABC} is nonnegative. While this statement is true for *any* quantum states, a special structure arises when the inequality is satisfied with an equality. In particular, Petz[17] showed that a tripartite state ρ_{ABC} can be reconstructed from ρ_{AB}, ρ_B and ρ_{BC} if the conditional mutual information vanishes:

$$\rho_{ABC} = \rho_{AB}^{\frac{1}{2}}\rho_B^{-\frac{1}{2}}\rho_{BC}\rho_B^{-\frac{1}{2}}\rho_{AB}^{\frac{1}{2}}. \tag{3}$$

Recursively applying Petz's result, Hastings and Poulin showed that quantum systems that saturate the equality condition for certain subsystems can be completely reconstructed from the local reduced density matrices alone.[14] A simple corollary of their result is that there cannot be another distinct state that is locally indistinguishable from the original state. To understand why, suppose there are two distinct states ρ_1, ρ_2 that are locally indistinguishable. Since both states have the same local reduced density matrices, the conditional mutual information computed from these local density matrices must be both 0. Due to the same reason, a recursive application of Eq.3 must yield a same recovered state for both ρ_1 and ρ_2. Since ρ_1 and ρ_2 were assumed to be distinct, this contradicts the original assumption. While this argument can be applied quite generally, it is not robust against small deviations of the original assumptions: if the conditional mutual information is only approximately equal to 0, Eq.3 cannot be used anymore. Also, Hastings and Poulin's result applies only to tree graphs. We shall discuss how these issues can be circumvented by giving several concrete examples.

–An inequality between topological entanglement entropy and topological degeneracy

We first discuss a class of models in two spatial dimensions that satisfy a strict form of area law. Loosely speaking, we say a model satisfies a strict form of area law if the nonuniversal contribution to the entanglement entropy can be canceled out by taking a judicious choice of a linear combination. The entanglement entropy of these models can be written as

$$S(A) = a|\partial A| - b_0(A)\gamma + O(e^{-|\partial A|/\xi}), \tag{4}$$

where a is a nonuniversal constant, $|\partial A|$ is the number of particles along the boundary of A, ξ is the correlation length, and γ is the topological entanglement entropy.[5, 6] $b_0(A)$ is the number of connected of components of the boundary of A. The correlation length will be equal to 0 for a fixed-point wavefunction of some renormalization group flow, but otherwise would remain nonzero.

Suppose we have N states $\{|\psi_i\rangle\}_{i=1,\cdots,N}$ with $(cL, O(e^{-L/\xi}))$-TQO condition that satisfy Eq.4 for a local subsystem. Here we have assumed our system to be on a $L\times L$ torus. The specific value of the numerical constant $0<c<1$ is irrelevant for the analysis, as long as it is sufficiently close to 1. Our first main result gives a rigorous lower bound of γ in terms of N, up to a small correction that vanishes in the thermodynamic limit.

$$\log N \leq 2\gamma + O(L^2 e^{-L/\xi}) \tag{5}$$

The idea for proving Eq.5 is to apply the Markov entropy decomposition(MED) to a maximally mixed state over the N states.[14] More precisely, consider a sequence of subsystems A_i, B_i, C_i, $i = 1, \cdots, n$ such that (i) $A_iB_iC_i = A_{i+1}B_{i+1}$, (ii) A_1B_1 and B_iC_i are *local*, and (iii) $A_nB_nC_n$ is the entire system. For such choice of subsystems, following linear combination of entanglement entropy is nonnegative due to the strong subadditivity of entropy:

$$\sum_{i=1}^{n} I(A_i : C_i|B_i) = S(A_1B_1) + \sum_{i=1}^{n} S(B_iC_i) - S(B_i) \\ - S(A_nB_nC_n).$$

By choosing the global state to be a uniform mixture of the N locally indistinguishable states, *i.e.* $\sum_{i=1}^{N} \frac{1}{N} |\psi_i\rangle \langle\psi_i|$, we arrive at the following bound:

$$\log N \leq S(A_1B_1) + \sum_{i=1}^{n} S(B_iC_i) - S(B_i). \tag{6}$$

Since A_1B_1, B_iC_i, and B_i are all local, their entanglement entropy can be replaced by an entanglement entropy of one of the states $|\psi_j\rangle$ with a small correction. The correction term can be estimated by using Fannes inequality[39] which holds for any quantum states ρ, σ supported on a d-dimensional Hilbert space:

$$|S(\rho) - S(\sigma)| \leq \epsilon \log d - \epsilon \log \epsilon, \quad \epsilon := |\rho - \sigma|_1, \tag{7}$$

where $|\cdots|_1$ is the trace norm. Eq.5 can be derived by choosing an appropriate set of subsystems such that the boundary contributions cancel out, while the universal term survives. One choice of such subsystems is depicted in FIG.1.

While we have assumed that the corrections to the area law and the local indistinguishability property decay exponentially in the system size, the same argument can be carried out even for an algebraically decaying function with a large enough exponent, see the Appendix. Eq.5 confirms the intuition that an amount of long-range entanglement limits the topological ground state degeneracy. For two-dimensional models supporting anyons, this result is already known. The ground state degeneracy of a topologically ordered system is n_p^2, where n_p is the number of particle types.[40] On the other hand, the topological entanglement entropy is related to the total quantum dimension of the system:

$$\gamma = \log \sqrt{\sum_a d_a^2},$$



FIG. 1: Each of the diagrams represent the subsystems A_i, B_i, C_i, $i = 1, 2, 3$ with the constraint that $A_iB_iC_i$ is equal to $A_{i+1}B_{i+1}$. Due to this construction, the entanglement entropy of $A_3B_3C_3$ is the only nonlocal contribution to the sum $\sum_{i=1}^{3} I(A_i : C_i|B_i)$. The entanglement entropy of $A_3B_3C_3$ becomes $\log N$ for a maximally mixed state over N locally indistinguishable states. The rest of the contributions can be computed from the formula for the entanglement entropy of local subsystems, *i.e.* Eq.4.

where d_a is the quantum dimension of a particle with a topological charge a.[5, 6] Since non-Abelian charges can have a quantum dimension that is strictly larger than 1, topological entanglement entropy has to be always larger or equal to $\frac{1}{2} \log N$.[5] However, we emphasize that we did not assume anything about the Hamiltonian at all. Eq.5 was derived only from the property of the states. Our result shows that even the topological ground state degeneracy - typically thought as a property inherited from the Hamiltonian - is already strongly constrained by the structure of the state alone. Admittedly our premise about the form of the entanglement entropy is quite restrictive. However, we note that there are number of models whose ground state entanglement entropy can be either analytically computed[5, 6, 25, 26, 32, 41] or computed with a good numerical precision[42–46].

– Higher dimensional systems

The preceding argument is based on the observation that the entanglement entropy of the subsystem can be decomposed into two parts. The first part is the local entanglement, which can be canceled out by making a judicious choice of subsystems. More precisely, we expect the entanglement entropy of a subsystem A to have the following form:

$$S(A) = S_{local}(A) + S_{topo}(A),$$

where $S_{local}(A)$ is an entanglement that can be decomposed into strictly local contributions.[47] The rest of the contributions, including the finite-size effects, are included in $S_{topo}(A)$. For such systems, the maximal number of topologically protected states depends on the scaling law of $S_{topo}(A)$. Alternatively, a quantum error-correcting code with an extensive ground state degeneracy would imply that there is an extensive subcorrection term for the entanglement entropy. Interesting examples include Chamon's model and Haah's cubic code, which are known to have ground state degeneracies that increase as $N = 2^{\Theta(L)}$ for certain choices of L.[48–50] Therefore, the entanglement entropy of these models must have a subcorrection term that grows as $\Omega(L)$.

Also, one may argue on a physical ground that the subcorrection term to the area law for gapped systems typically scales as $|A|^{D-2}$.[47] For such systems, the maximal number of topologically protected qubits is bounded by $O(L^{D-2})$.

– *Bounds for more generic systems*

In general, one cannot expect the leading terms of the entanglement entropy to be canceled out by choosing an appropriate set of subsystems, especially for critical systems and ground states of nonlocal Hamiltonian. We show that, even for such generic systems, a nontrivial tradeoff bound can be obtained. Using the subadditivity of entropy, *i.e.* $I(A:B) = S(A) + S(B) - S(AB) \geq 0$, Eq.6 yields the following inequality:

$$k \leq \sum_i S(X_i), \quad |X_i| < d \tag{8}$$

for a quantum code with a code distance d and a number of encoded qubits $k = \log N$, where $\{X_i\}$ is a partition of the system. Dividing both sides of the inequality by the number of particles n, we find that the rate of a quantum error correcting code $\frac{k}{n}$ is bounded by the average entanglement entropy per volume over any partitions $\{X_i\}$, $|X_i| < d$. Our result shows that studying the entanglement properties of a quantum error correcting code is a relevant problem for understanding its fundamental limit. In particular, Eq.8 gives a necessary condition for a quantum error correcting code to have a nonvanishing rate: its average entanglement entropy over subsystems smaller than the code distance must satisfy a strict volume law.

Eq.8 can be used to obtain a tradeoff bound for quantum codes satisfying a subvolume law of entanglement entropy. If the entanglement entropy satisfies a subvolume law,

$$S(A) = O(|A|^{\alpha}),$$

we can obtain the following tradeoff bound:

$$kd^{1-\alpha} = O(n). \tag{9}$$

There are at least two important differences between Eq.1 and Eq.9. On one hand, Eq.9 is more general than Eq.1 in that it does not require any structure about the parent Hamiltonian. On the other hand, Eq.9 provides a weaker tradeoff bound than Eq.1 does.

– *Outlook*

MED provides a powerful numerical framework for studying finite-temperature quantum many-body systems.[14, 51] By extending the observation of Hastings and Poulin, we have shown that MED can be also used to obtain a fundamental limit on the information storage capacity of a quantum many-body system. In particular, we have shown that there is a fundamental reason behind the existence of the topological entanglement entropy. After all, a state with a sufficiently small topological entanglement entropy cannot have another state that is locally indistinguishable from the original state. It is quite remarkable that such a tight tradeoff bound can be obtained simply from the property of the wavefunction alone. There are several interesting physical systems for which our approach might be useful.

For example, our approach should be easily generalizable to finite temperature systems. In such systems, the topological entanglement entropy is interpreted as an order parameter characterizing the phase.[52–55] While the technical details must be worked out, we can have an intuitive explanation on why those terms must be $\Omega(1)$. Some of these models become a self-correcting classical/quantum memory depending on the temperature.[56, 57] Since the "logical operators" that are associated to these systems are highly nonlocal, two distinct states cannot be distinguished via any local operation. If the topological entanglement entropy becomes sufficiently small, the number of locally indistinguishable states is bounded by a small number, which would give rise to a contradiction. It is well-known that Eq.4 is modified if the subsystem contains a quasiparticle with a nontrivial topological charge.[5] It would be interesting to extend our analysis to such settings as well.

We conclude with a remark that our result can be used as a rigorous tool for proving a distinctiveness of different quantum many-body phases. Given a quantum many-body system on a torus with N topologically protected ground states, it cannot be adiabatically connected to *any* state with a topological entanglement entropy strictly lower than $\frac{\log N}{2}$. This is due to the fact that topologically protected states remain to be so under an adiabatic evolution.[37, 58] Therefore, the lower bound for the topological entanglement entropy, *i.e.* Eq.5, remains stable with a small correction that vanishes in the thermodynamic limit.

– *Acknowledgements*

I would like to thank John Preskill, Alioscia Hamma, and Roger Mong for helpful discussions and suggestions. Part of this work was done while the author was attending 2013 Coogee quantum information theory workshop. This research was supported in part by NSF under Grant No. PHY-0803371, by ARO Grant No. W911NF-09-1-0442, and DOE Grant No. DE-FG03-92-ER40701.

---

[1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Rev.Mod.Phys. **81**, 865 (2009), quant-ph/0702225.

[2] J. Eisert, M. Cramer, and M. B. Plenio, Rev. Mod. Phys. **82,**, 277 (2008), 0808.3773.

[3] P. Calabrese and J. Cardy, J.Stat.Mech. p. P06002 (2004), hep-th/0405152.

[4] A. Hamma, R. Ionicioiu, and P. Zanardi, Phys.Lett. A **337**, 22 (2005), quant-ph/0406202.

[5] A. Kitaev and J. Preskill, Phys.Rev.Lett. **96**, 110404 (2006), hep-th/0510092.
[6] M. Levin and X.-G. Wen, Phys. Rev. Lett. **96**, 110405 (2006), cond-mat/0510613.
[7] B. N. M. Fannes and R. F. Werner, Comm. Math. Phys. **144**, 443 (1992).
[8] D. Perez-Garcia, F. Verstraete, M. M. Wolf, and J. I. Cirac, Quantum Inf. Comput. **7,**, 401 (2007), quant-ph/0608197.
[9] F. Verstraete and J. I. Cirac (2007), cond-mat/0407066.
[10] G. Vidal, Phys. Rev. Lett. **101**, 110501 (2008), quant-ph/0610099.
[11] M. B. Hastings, JSTAT, P **08024** (2007), 0705.2024.
[12] F. G. S. L. Brandao and M. Horodecki (2012), 1206.2947.
[13] I. Arad, A. Kitaev, Z. Landau, and U. Vazirani (2013), 1301.1162.
[14] D. Poulin and M. B. Hastings, Phys. Rev. Lett. **106**, 080403 (2010), 1012.2050.
[15] I. H. Kim (2012), 1210.1831.
[16] I. H. Kim, Phys. Rev. B **86,**, 245116 (2012), 1210.2360.
[17] D. Petz, Rev. Math. Phys. **15**, 79 (2003), quant-ph/0209053.
[18] P. Hayden, R. Jozsa, D. Petz, and A. Winter, Commun. Math. Phys. **246**, 359 (2004).
[19] M. Leifer and D. Poulin, Ann. Phys. **323**, 1899 (2007), 0708.1337.
[20] S. Bravyi and B. Terhal, New J. Phys. **11**, 043029 (2008), 0810.1983.
[21] S. Bravyi, D. Poulin, and B. Terhal, Phys. Rev. Lett. **104**, 050503 (2009), 0909.5200.
[22] J. Haah and J. Preskill, Phys. Rev. A **86,**, 032308 (2010), 1011.3529.
[23] O. Landon-Cardinal and D. Poulin (2012), 1209.5750.
[24] N. Delfosse (2013), 1301.6588.
[25] A. Y. Kitaev, Annals Phys. **303**, 2 (2003), quant-ph/9707021.
[26] M. A. Levin and X.-G. Wen, Phys.Rev. B **71**, 045110 (2005), cond-mat/0404617.
[27] B. I. Halperin, Phys. Rev. B **25**, 2185 (1982).
[28] X.-G. Wen, Int. J. Mod. Phys. B **6**, 1711 (1992).
[29] R. Moessner and S. L. Sondhi, Phys. Rev. Lett **86**, 1881 (2001), cond-mat/0007378.
[30] R. Moessner and S. L. Sondhi, Phys. Rev. B **68**, 184512 (2003), cond-mat/0307592.
[31] D. Perez-Garcia, F. Verstraete, J. I. Cirac, and M. M. Wolf, Quant. Inf. Comp. **8**, 0650 (2007), 0707.2260.
[32] N. Schuch, I. Cirac, and D. Perez-Garcia, Annals of Physics **325**, 2153 (2010), 1001.3807.
[33] A. Kitaev, Annals of Physics **321**, 2 (2006), cond-mat/0506438.
[34] E. Knill and R. Laflamme, Phys. Rev. A **55**, 900 (1997).
[35] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information*, Cambridge Series on Information and the Natural Sciences (Cambridge University Press, 2000), ISBN 9780521635035.
[36] D. W. Kribs, R. Laflamme, D. Poulin, and M. Lesosky, Quant. Inf. & Comp. **6**, 383 (2006), quant-ph/0504189.
[37] S. Bravyi, M. B. Hastings, and F. Verstraete, Phys. Rev. Lett. **97**, 050401 (2006), quant-ph/0603121.
[38] E. H. Lieb and M. B. Ruskai, J. Math. Phys. **14**, 1938 (1973).
[39] M. Fannes, Comm. Math. Phys. **31**, 291 (1973).
[40] X. G. Wen and Q. Niu, Phys. Rev. B **41**, 9377 (1990).
[41] H. Yao and X.-L. Qi, Phys. Rev. Lett., **105,**, 080501 (2010), 1001.1165.
[42] M. Haque, O. Zozulya, and K. Schoutens, Phys. Rev. Lett. **98**, 060401 (2007).
[43] O. S. Zozulya, M. Haque, K. Schoutens, and E. H. Rezayi, Phys. Rev. B **76**, 125310 (2007).
[44] S. Furukawa and G. Misguich, Phys. Rev. B **75**, 214407 (2007).
[45] Y. Zhang, T. Grover, and A. Vishwanath, Phys.Rev.B **84**, 075128 (2011), 1106.0015.
[46] D. Poilblanc, N. Schuch, D. Pérez-García, and J. I. Cirac, Phys. Rev. B **86**, 014404 (2012), 1202.0947.
[47] T. Grover, A. M. Turner, and A. Vishwanath, Phys. Rev. B **84**, 195120 (2011), 1108.4038.
[48] C. Chamon, Phys. Rev. Lett. **94**, 040402 (2005).
[49] S. Bravyi, B. Leemhuis, and B. M. Terhal, Ann. Phys. **326**, 839 (2010), 1006.4871.
[50] J. Haah, Phys. Rev. A **83**, 042330 (2011).
[51] A. J. Ferris and D. Poulin (2012), 1212.1442.
[52] C. Castelnovo and C. Chamon, Phys. Rev. B **76**, 184442 (2007), 0704.3616.
[53] C. Castelnovo and C. Chamon, Phys. Rev. B **78**, 155120 (2008), 0804.3591.
[54] S. Iblisdir, D. Perez-Garcia, M. Aguado, and J. Pachos, Nuclear Physics B **829**, 401 (2010).
[55] D. Mazac and A. Hamma, Ann. Phys. **327**, 2096 (2011), 1112.0947.
[56] R. Alicki, M. Horodecki, P. Horodecki, and R. Horodecki, Open Syst. Inf. Dyn. **17**, 1 (2008), 0811.0033.
[57] S. Chesi, D. Loss, S. Bravyi, and B. M. Terhal, New J. Phys. **12,**, 025013 (2009), 0907.2807.
[58] S. Bravyi, M. Hastings, and S. Michalakis, J. Math. Phys. **51**, 093512 (2010), 1001.0344.
[59] We note in passing that classical error correcting codes in general do not satisfy Eq.2, *e.g.* a classical repetition code. Therefore, the same argument cannot be applied to such codes.

Derivation of Eq.5 under a power law correction

In the main text of this Letter, we have derived an inequality between the topological entanglement entropy γ and the ground state degeneracy N on a torus. There are several assumptions that we have implicitly assumed in the analysis. First, we have assumed that the entanglement entropy of a region that can be contained in a radius of cL for a sufficiently large value of $0 < c < 1$ is expressed as Eq.4. However, we did not specify the value of c explicitly. We have also assumed that the corrections to the area law and the local indistinguishability property decays exponentially in the system size. Here we show that these conditions can be relaxed significantly in general.

More precisely, we obtain an inequality analogous to Eq.5 under the following set of assumptions.

- The system satisfies a (cL, ϵ)-TQO condition with $0 < c < 1$, with $\epsilon = O(\frac{1}{L^{\alpha_1}})$, $\alpha_1 > 0$.
- The number of locally indistinguishable states satisfying the TQO condition is N.

- The correction to the area law decays algebraically, *i.e.* $S(A) = a|\partial A| - \gamma + O(\frac{1}{|A|^{\alpha_2}})$, $\alpha_2 > 0$.
- The Hilbert space dimension of the local particles is d.

We begin by bounding a difference between the entanglement entropy of a single wavefunction, say $|\psi_1\rangle$, and the entanglement entropy of a mixed state $\rho = \frac{1}{N}\sum_{i=1}^{N} |\psi_i\rangle\langle\psi_i|$. Without loss of generality, we shall assume that Eq.4 is the entanglement entropy formula for $|\psi_1\rangle$. Denoting the entanglement entropy of the single wavefunction as $S^1(A)$ and the entanglement entropy of the mixed state ρ as $S^\rho(A)$, we obtain the following bound.

Lemma 1. *Let A be a region that can be contained in a ball of radius cL.*

$$|S^1(A) - S^\rho(A)| \leq \epsilon(d|A| - \log\epsilon) \tag{10}$$

This result follows from a simple application of the Fannes inequality, see Eq.7.

As we did in the main text, consider a sequence of subsystems A_i, B_i, and C_i such that $A_iB_iC_i = A_{i+1}B_{i+1}$. By choosing the subsystems A_1B_1 and B_iC_i to be small enough to be contained in a ball of radius cL,

$$\log N \leq S^\rho(A_1B_1) + \sum_{i=1}^{n} S^\rho(B_iC_i) - S^\rho(B_i) \tag{11}$$

$$\leq S^1(A_1B_1) + \sum_{i=1}^{n} S^1(B_iC_i) - S^1(B_i) + \epsilon_{LA}, \tag{12}$$

where ϵ_{LA} is the approximation error from Eq.10. One can choose the size of the subsystems A_1B_1 and B_iC_i to be $\Theta(c^2L^2)$, in which case n becomes $n = \Theta(\frac{1}{c^2})$. Therefore,

$$\epsilon_{LA} = O(\epsilon d L^2). \tag{13}$$

Also,

$$S^1(A_1B_1) + \sum_{i=1}^{n} S^1(B_iC_i) - S^1(B_i) = 2\gamma + O(\frac{c^{-2(1+\alpha_2)}}{L^{2\alpha_2}}). \tag{14}$$

Combining these bounds together, for a constant c and d, we conclude that

$$\log N \leq 2\gamma + O(L^{2-\alpha_1}) + O(L^{-2\alpha_2}). \tag{15}$$

Therefore, for a sufficiently large system size, our main result holds for any $\alpha_1 > 2$, $\alpha_2 > 0$. The significance of this bound comes from the fact that a conservative estimate for the area law correction term gives $\alpha_2 = \frac{1}{2}$.[5, 47] The preceding analysis shows the robustness of our bound even under such a conservative error estimate. Even under an algebraic correction to the ideal wavefunction, the inequality remains intact.

We would also like to explain the intuition behind why a factor of 2 appears in front of γ. It should be clear from the construction that the area terms cancel out. Therefore, one only needs to be concerned about the topology of the subsystems A_1B_1, B_iC_i, and B_i. The general idea is that for the construction in FIG.1, A_1B_1 and B_1C_1 are simply connected, yet B_1 is a union of two simply connected subsystems. Hence the topological contribution must be γ for A_1B_1 and B_1C_1, and 2γ for B_1. Similarly, B_2C_2 gives a contribution of γ, while B_2 gives 2γ. B_3C_3 is simply connected, so it gives a contribution of γ. B_3 is an annulus, hence it gives 2γ. Combining these results together, the topological contribution to our bound becomes 2γ. A similar analysis can be carried out for other choice of subsystems as well. What is important is that Eq.11 holds for *any* choice of subsystems. Therefore, one can always optimize over different set of subsystems to obtain the best bound.